GIBSON COUNTY: Key Data Points

- Gibson county experiences a slightly lower total birth rate than the state of Indiana overall, 13 versus 13.9 respectively.
- The county's mothers are giving birth to a higher percentage of low birth weight babies when compared to state, 8.1% versus 7.9% respectively.
- Compared to the state as a whole, Gibson county has a larger percentage of mothers smoking during pregnancy, 23.5% versus 18.5% respectively.
- A much lower percentage of Gibson county mothers breastfeed on discharge, compared to the state (50.9% vs. 63.2%, respectively)
- Gibson county observed 20 accidental deaths (10 of which were motor vehicle), 5 suicidal deaths and 2 homicide

Note: All figures relate to 2003 data from the Indiana State Department of Health, unless otherwise specified

Figure 1. Births

	Total		White		Black		Other	
	Number	Crude Rate*	Number	Crude Rate*	Number	Crude Rate*	Number	Crude Rate*
Gibson County	430	13	417	13.1	6	9.2(U)	7	15.1(U)
INDIANA	86,382	13.9	75,422	13.7	9,288	17.5	1,672	10.6

^{*} births per 1,000 population; U = unstable rate, based on fewer than 20 births

Figure 2. Mortality

		NUMBER				RATE*		
COUNTY OF	REPORTED RACE/	Live	Infant Deaths (< 1 yr)	Neonatal Deaths (< 28 days)	Post Neonatal Deaths	Infant Deaths (< 1 yr)	Neonatal Deaths (< 28 days)	Post Neonatal Deaths
RESIDENCE	ETHNICITY	Births			(28-364 days)			(28-364 days)
GIBSON	All Races	430	1	1	0	**	**	0
	White	417	1	1	0	**	**	0
	Black	6	0	0	0	0	0	0
	Other	7	0	0	0	0	0	0
	Hispanic	2	0	0	0	0	0	0
INDIANA	All Races	86,382	641	417	224	7.4	4.8	2.6
	White	75,422	479	309	170	6.4	4.1	2.3
	Black	9,288	148	98	50	15.9	10.6	5.4
	Other	1,672	14	10	4	**	**	**
	Hispanic	6,763	38	22	16	5.6	3.3	**

^{*}rate per 1,000 live births **Numerator is less than 20 and the rate is unstable

Figure 3. Outcome Indicators of Mother, by Race


		Birth W	/eight			During P	regnancy	
GIBSON	Total	% Low	% Very Low	% Preterm	% PNC 1st	%	%	% Unmarried
COUNTY	Births	(< 2500 grams)	(< 1500 grams)	†	Trimester [‡]	Alcohol	Smoked	Parents
All Races	430	8.1	0.9	8.8	86.3	1.2	23.5(S)	29.8
White	417	8.2	1	8.9	85.9	1.2	23.3	28.5
Black	6	**	**	**	**	**	**	**

Birth Weight			/eight			During P	regnancy	
	Total	% Low	% Very Low	% Preterm	% PNC 1st	%	%	% Unmarried
INDIANA	Births	(< 2500 grams)	(< 1500 grams)	†	Trimester ‡	Alcohol	Smoked	Parents
All Races	86,382	7.9	1.4	9.9	80.6	0.6	18.5	37.1
White	75,422	7.2	1.2	9.5	82.1	0.6	19.1	32.6
Black	9,288	13.3	3.3	13.3	68.2	1.1	15.2	76.3

[†] Preterm: less than 37 weeks gestation

Figure 4. Breastfeeding

GIBSON COUNTY	All Races	White	Black
Total Births	430	417	6
Breastfeeding on Discharge	219	212	3
% Breastfeeding on Discharge	50.9	50.8	50



[†] PNC: Prenatal Care

'S' Significantly different than the state percent

"**' Outcomes for this county are suppressed due to 20 or fewer total births

Figure 4. Breastfeeding

INDIANA	All Races	White	Black
Total Births	86,382	75,422	9,288
Breastfeeding on Discharge	54,636	49,385	3,959
% Breastfeeding on Discharge	63.2	65.5	42.6




Figure 5. HIV/AIDS

	Total # of AIDS cases reported since 1982	Total # of HIV cases reported since 1985
GIBSON COUNTY	13	4
INDIANA	7,641	3,856

^{*}data gathered from June 2005 HIV/STD Quarterly Report

Figure 6. Nutritional Risk Factors by Status,

Indiana WIC Clinics, Quarter Ending 09/30/2004

Local Agency: Gibson County Rehabilitation Centers, Inc.								
Counties Included: Gibson, Posey								
	Childr	en 1-5 y	ears of Ag	e			%	
Risk Factor or Status	White	Black	Hispanic	Asian	Other	Total	within Agency	
Overweight	25	1	1	1	1	29	6.2%	
At Risk for becoming Overweight	22	2	0	0	2	26	5.6%	

Local Agency: State Wide Counties Included: All

Countres meruded. 7m								
	Childr	en 1-5 y		%				
Risk Factor or Status	White	Black	Hispanic	Asian	Other	Total	within Agency	
Overweight	1,732	445	800	12	97	3,086	4.8%	
At Risk for becoming Overweight	2,023	485	639	11	124	3,282	5.1%	

Priority Issues (Gibson & Indiana	a)	
Demographics: Percentage of population enrolled in Medicaid, Percentage of population enrolled in Food Stamps.	Gibson	Indiana
2 % of Population enrolled in Medicaid as of June 2003	10.3	12.4
3 % of Population in Food Stamps Program, 2004	5.4	8.6
Priority #1 (pregnancy outcomes): Average Infant Mortality Rate, Average Fetal death Rate	Gibson	Indiana
1 Av Total Infant Death Rate (1993-2002)	7.5	8.1
2 Average Fetal Death Rate (1994-2003)	7.0	6.7
Priority #2 (reduce barriers): Health Professional Shortage Areas, Percentage of women with prenatal care in first trimester	Gibson	Indiana
Scores for Primary Care Health Professional Shortage Areas (HPSAs) (2005)	No Score	NA
2 % PNC 1st Trimester†, 2003	86.3	80.6
Priority #4 (environmental): Prevalence of asthma, Childhood lead poisoning	Gibson	Indiana
Prevalence of Asthma in a Medicaid Cohort of Children age 0-17 years, 2003	7.1	10.4
2 %Estimated Lead Blood Levels (%EBLL) for Children under 6 years old FY 2004-2005	4.3	2.1
Priority #5 (tobacco): Percentage of mothers who smoked during pregnancy, Percentage of Adult Smoking	Gibson	Indiana
1 % of mothers Smoked During Pregnancy, 2003	23.5	18.5
2 % of Adult Smoking in 2003	31.6	29.3
Priority #7 (adolescent risk behaviors): Adolescent deaths due to motor vehicle accidents, Chlamydia rates (ages 15-24).	Gibson	Indiana
Number of Adolescent (15-24) deaths due to Motor vehicle accidents (2001-2003)	6	748
2 Chlamydia rates per 100,000 county specific population (15-24yrs of Age)	610.15	1541.49
Priority #8 (obesity): Obesity in pregnant women	Gibson	Indiana
Number of Obese Pregnant Women In Indiana, 2003	20	2,496
Priority #9 (domestic violence, childhood injury): Child deaths due to Injury, Child abuse and neglect rates	Gibson	Indiana
1 Child deaths due to Injury 1-14yrs (2001-2003)	1	257
2 Child Abuse and Neglect per 1,000 Population under age 18, 2004	16.3	13.5
Priority #10 (To improve racial and ethnic disparities in women of childbearing age, mothers, and children's health outcomes): Percentage of Hispanic Population of any Race, LBW Disparity Ratio	Gibson	Indiana
1 % of Hispanic Population of any Race, July,1,2004	0.8	4.3
2 Low Birth Weight Disparity Ratio (B/W) (1993-2002)	NS	1.92

Source: Data Analysis Team, ISDH. **Notes:** 1. NS= Not Stable 2. NA = Not Available 3. Infant Mortality rate and Fetal Death rate is for 1000 live births. 4. Low Birth Weight: Birth Weight Less than 2,500grams. 5. HPSA Score ranges 1 – 25. The higher the score, the higher the need. 6. EBLL: Ten (10) micrograms of lead per deciliter of whole blood for one venous test; %EBLL is based on screening rate of 27% Statewide, unduplicated kids 7. Obesity in Pregnant Women determined by "Concurrent Illness"

Obesity" checkbox on the birth certificate 8. PNC percentages are calculated using total births in each county.